

# REFLEXES OF SPINAL AND CRANIAL NERVES

**SENSORY  
STIMULUS**



**MOTOR  
RESPONSE**


**Definition of a Reflex - stereotyped motor response to a specific sensory stimulus**

**AFFERENT ARM  
OF REFLEX  
SENSORY  
STIMULUS**

## **TYPICAL REFLEX**

**EFFERENT ARM  
OF REFLEX  
MOTOR  
RESPONSE**

**in Central Nervous System**



- Typical reflex arc: 1) sensory neuron - detects stimulus (termed afferent arm of reflex arc)**
- 2) interneurons - (most often) effects on motor neuron can be excitatory or inhibitory**
- 3) motor neurons - produce muscle contraction, motor response (termed efferent arm of reflex arc)**

**For reflex to occur, all elements must be functional:  
If absent, diagnose where pathway is interrupted.  
If abnormal, diagnose where pathway is compromised.**

**REFLEXES CAN BE USED TO TEST NERVOUS SYSTEM FUNCTION,  
LOCATE SITE OF LESION**

# EVALUATING REFLEXES

**TABLE 21-8** Scoring Deep Tendon Reflexes

Grade	Deep Tendon Reflex Response
0	No response
1+	Sluggish or diminished
2+	Active or expected response
3+	More brisk than expected, slightly hyperactive
4+	Brisk, hyperactive, with intermittent or transient clonus

**NOTE: DEEP TENDON REFLEX = STRETCH REFLEX**


**Reflex is evaluated according to:**

- 1) amount (size, magnitude) of motor response,**
- 2) latency (time to elicit motor response);**

**Hyper-reflexia = enhanced reflexes; in some disease processes, damage can enhance reflex responses**


**Clonus = series of abnormal, rapid alternating contractions and relaxations of muscle produced by single stimulus**

## SOME REFLEXES ARE PROTECTIVE AND CONSTANT



Ex. Pupillary light reflex – shine light in eye, pupil constricts

## SOME REFLEXES ARE CONSTANT UNDER SAME CIRCUMSTANCES



STRETCH (DEEP TENDON)  
REFLEXES - can be tested in a number of muscles; activate muscle spindles

- 1) Patient positioned correctly, told to relax; focus patient's attention elsewhere (ex. tell patient to clench hands and try pulling apart);
- 2) COMPARE REFLEXES ON RIGHT AND LEFT SIDES - Reason: reflexes can be modulated (changed or blocked) by activities in CNS.

## II. SPINAL REFLEXES

Three basic reflexes:

A) Stretch reflex - produced by activating Muscle Spindles - contributes to maintaining postural stability, countering sudden loads


B) Autogenic inhibition - produced by activating Golgi tendon organs - aids in regulating muscle tension, prevents damage to tendon, bone

C) Flexion reflex - produced by activating Cutaneous, Pain receptors - avoid obstacle or painful stimulus (stepping on nail)

# REFLEXES CAN HAVE WIDESPREAD EFFECTS


## EFFECTS:

1. Excite muscle containing the sense organ;
2. Excite muscles with same action (termed Synergist muscles)
3. Inhibit muscles with opposite action (termed Antagonist)
4. Some reflexes have effects in opposite limb (termed Contralateral muscles; ex. in flexor reflex, reflex causes lifting of one leg and extension in opposite leg to prevent falling).



## STIMULUS


**BICEPS  
MUSCLE  
STRETCHED**



**1) Stimulus -  
fast stretch  
of muscle**

## A. STRETCH REFLEX

**BICEPS MUSCLE SPINDLE**




**2) Sense organ  
excited - Muscle  
spindle Ia and II  
sensory neurons**

## RESPONSE




**BICEPS  
MUSCLE  
CONTRACTS**



**3) Primary  
response -  
muscle that is  
stretched  
contracts rapidly**


# SENSORY EXCITED - MUSCLE SPINDLES - SPINDLES HAVE SENSORY AND MOTOR (GAMMA) NEURONS



**Spindle muscle cells are specialized skeletal muscle cells innervated both by Sensory neurons (cell bodies in dorsal root ganglia) and Motor neurons (cell bodies in ventral horn)**  
**MUSCLE SPINDLES ARE STRETCHED AND SENSORY NEURONS DISCHARGE WHEN A MUSCLE IS STRETCHED**



# MUSCLE SPINDLES HAVE BOTH SENSORY AND MOTOR INNERVATION (GAMMA MOTOR NEURONS);



Gamma motor neurons innervate only muscle cells in muscle spindles, not regular skeletal muscle cells (alpha motor neurons); Gamma motor neurons can adjust sensitivity of muscle spindles to stretch (nervous patient shows exaggerated stretch reflexes).

# GAMMA DYNAMIC MOTOR NEURONS CAN ENHANCE SENSITIVITIES OF SPINDLE SENSORY NEURONS




Gamma motor neuron activity is increased in anticipation of perturbations (ex. walking on a thin rope)



Gamma motor neuron activity is increased (probably) in patients who are nervous.

## STIMULUS



**BICEPS  
MUSCLE  
STRETCHED**



**1) Stimulus -  
fast stretch  
of muscle**


## A. STRETCH REFLEX

**BICEPS MUSCLE SPINDLE**




**2) Sense organ  
excited - Muscle  
spindle Ia and II  
sensory neurons**

## RESPONSE




**BICEPS  
MUSCLE  
CONTRACTS**




**3) Primary  
response -  
muscle that is  
stretched  
contracts rapidly**

Ia




# EFFECT ON MUSCLE: MONOSYNAPTIC CONNECTION



**Group Ia** - signal movements (rate of stretch) - **monosynaptic** connections with alpha motor neurons (**fastest reflex known, delay at synapse about 1 msec**)

**Group II** – signal position (amount of stretch) response weaker make 1) **monosynaptic** and 2) **polysynaptic** (through interneuron)


**+** note: **plus** indicates **excitatory** connection



# OTHER COMPONENTS OF STRETCH REFLEX




1) Excite synergist muscles - spindle afferents also make excitatory **monosynaptic** connections with synergist muscles



2) Inhibit antagonist muscles - **RECIPROCAL INHIBITION** - Spindle activity also excites **interneurons** that make **inhibitory synapses** on motor neurons to antagonist muscles (**polysynaptic**)

SENSE ORGAN =  
Biceps  
Muscle  
Spindle




# MUSCLE TONUS

- 1- Because connection is monosynaptic, ongoing activity in muscle spindles is important in determining firing of alpha motor neurons at rest.
- 2- Eliminating activity of spindles can decrease motor neuron firing producing decreased tonus.
- 3- Increased sensory activity can increase tonus.

## CLINICAL TESTING OF STRETCH REFLEX: TENDON TAP


- 1- Tendon tap elicits twitch because it **excites almost all muscle spindles simultaneously**
- 2- **Excitation converges upon motor neuron**



**Spasticity/Rigidity – Increased tonus occurs after Upper Motor Neuron Lesion (ex. stroke); due to loss of modulation of reflex**

## STIMULUS



**BICEPS  
MUSCLE  
STRETCHED**



**1) Stimulus -  
fast stretch  
of muscle**


# A. STRETCH REFLEX

**BICEPS MUSCLE SPINDLE**



**2) Sense organ  
excited - Muscle  
spindle Ia and II  
sensory neurons**


Ia



## RESPONSE



**BICEPS  
MUSCLE  
CONTRACTS**



**3) Primary  
response -  
muscle that is  
stretched  
contracts rapidly**


# CLINICAL TESTING OF STRETCH REFLEX: TENDON TAP

**NOTE: COMPARE REFLEXES ON RIGHT AND LEFT SIDES**






# REFLEXES MUST BE MODIFIED DURING VOLUNTARY MOVEMENTS



Voluntary contraction of one muscle often produces stretch of the antagonist muscle. If stretch reflexes were always active, voluntary contraction of one muscle would produce reflex contraction in the antagonist.

- Therefore, stretch reflexes can be modified in some muscles during voluntary movements

# MODIFICATION OF REFLEXES: MECHANISMS





1- Reflexes can be modulated by

- 1) Gamma motor neurons – change muscle spindle sensitivity
- 2) Descending inputs from brain – some produce pre-synaptic inhibition of Ia terminals; some change excitability of motor neurons..

**Changes in reflexes are symptomatic: In general, Decreased Stretch reflexes can indicate Lower Motor Neuron Disorders, Increased Stretch reflexes can indicate Upper Motor Neuron Syndromes.**

# REMINDER: UPPER VS LOWER MOTOR NEURON




# HYPERREFLEXIA: INCREASED STRETCH REFLEX ON ONE SIDE

[used by permission of Paul D. Larsen, M.D., University of Nebraska Medical Center;  
<http://library.med.utah.edu/neurologicexam>]



# Camillo Golgi (1843-1926)

excellent  
mustache




1) Stimulus -  
Large force  
exerted on  
muscle tendon

2) Sense organ  
excited -  
Golgi tendon  
organs - located in  
muscle tendon,  
signal FORCE


## B. AUTOGENIC INHIBITION

MUSCLE  
TENSION  
INHIBITED



3) Primary response -  
muscle  
attached to  
tendon relaxes

# AUTOGENIC INHIBITION




## PRIMARY RESPONSE

Synapses - polysynaptic

- 1) Ib sensory neuron (GTO) makes excitatory synapse onto interneuron
- 2) Interneuron makes inhibitory synapse onto motor neuron; Motor neuron decreases firing

Function of Autogenic inhibition - Regulating muscle tensions  
(protective, prevent damage to tendon)



# AUTOGENIC INHIBITION

## Other effects


### a. Inhibits synergist muscles -

GTO makes excitatory synapse on interneuron; interneuron makes inhibitory synapse on motor neurons to synergist muscle

### b. Excites antagonist muscles -



GTO makes excitatory synapse on interneuron; interneuron makes excitatory synapse on motor neurons to antagonist muscles

**CLASPED KNIFE REFLEX:** in Upper motor neuron lesions, tonus increases, resistance to stretch increases; if sufficient force is applied, limb resistance suddenly decreases (like pocket knife snapping shut)



**CLASPED KNIFE REFLEX:** is an example of Autogenic inhibition. It is elicited in patients with UMN lesions due to high tonus in muscle.


1) PHYSICIAN TRIES TO FLEX ELBOW JOINT OF PATIENT WITH UPPER MOTOR NEURON LESION




PHYSICIAN HOLDS WRIST AND PUSHES HERE AFTER TELLING PATIENT TO RELAX

ENCOUNTERS HIGH RESISTANCE DUE TO HIGH TONUS IN TRICEPS AND HIGH STRETCH REFLEXES


2) KEEP TRYING AND TENSION ON TRICEPS TENDON EXCITES GOLGI TENDON ORGANS



HIGH IMPOSED FORCE EXCITES GOLGI TENDON ORGANS IN TRICEPS TENDON WHICH INHIBITS MOTOR NEURONS TO TRICEPS MUSCLE



3) TRICEPS RELAXES AND RESISTANCE SUDDENLY DECREASES: ELBOW JOINT FLEXES



ELBOW JOINT SNAPS SHUT LIKE A POCKET KNIFE = CLASPED KNIFE REFLEX



# AUTOGENIC INHIBITION AND FORCE REGULATION

1- **Regulating muscle tension** - forces developed by contractions of muscles are automatically controlled so that they **do not cause damage to tendons** (ex. lifting heavy object).




2- **Regulation of force during other behavior is more complex** (ex. walking) –

**Connections for autogenic inhibition may be inactivated during walking**

**Effects of Golgi tendon organs can then become excitatory via other interneurons**



# C. FLEXOR REFLEX




1) Stimulus - painful or noxious stimulus (stepping on nail)

2) Sense organ excited - Cutaneous receptors, Pain receptors (nociceptors)

3) Primary response - Protective withdrawal of limb


# FLEXOR REFLEX: PATHWAYS



## Synapses - Polysynaptic

1) Cutaneous afferent makes excitatory synapse onto Interneuron; Interneuron can synapse upon another interneuron

2) Interneuron makes excitatory synapse onto Flexor motor neuron





# FLEXOR REFLEX: OTHER EFFECTS

## ALL ARE POLYSYNPACTIC BY INTERNEURONS

1) Excite synergist muscles - excite other flexors in same leg (other joints)


2) Inhibit antagonist muscles - inhibit Extensors in same leg

3) **CROSSED EXTENSION REFLEX - EXCITE EXTENSORS AND INHIBIT FLEXORS IN OPPOSITE LEG**




**FUNCTION: OTHER LEG PROVIDES SUPPORT WHEN FIRST LEG IS LIFTED**

# REFLEXES ARE MODULATED: SOME FLEXOR REFLEXES CAN CHANGE AFTER LESIONS, DISEASE PROCESSES



**NORMAL RESPONSE**


**STIMULUS –  
TO SKIN  
OF SOLE  
OF FOOT**



**FLEX  
TOES  
(DOWN)**



**BABINSKI SIGN –  
(EXTENSOR PLANTAR  
RESPONSE)**



**EXTEND BIG  
TOE, FANNING  
(ABDUCTION)  
OF OTHER  
TOES**

**Babinski sign** - seen after **Upper Motor neuron lesion**  
-direction of movement **changes from flexing toes to  
extending and fanning (abducting) toes**

**PLANTAR REFLEX: 'FLEXOR' REFLEX (PLANTAR FLEXION) IN FOOT: NORMAL** [used by permission of Paul D. Larsen, M.D., University of Nebraska Medical Center; <http://library.med.utah.edu/neurologicexam>]



**PLANTAR REFLEX: ABNORMAL, (POSITIVE) BABINSKI SIGN ON ONE SIDE** [used by permission of Paul D. Larsen, M.D., University of Nebraska Medical Center; <http://library.med.utah.edu/neurologicexam>]



# **\*\* 1. PUPILLARY LIGHT REFLEX - II TO III**

**AFFERENT ARM OF REFLEX**

**SENSORY  
STIMULUS**

**LIGHT IN  
EYE**



**EFFERENT ARM OF REFLEX**

**MOTOR  
RESPONSE**


**CONSTRICT  
PUPIL**






# PUPILLARY LIGHT REFLEX

**CN II - OPTIC NERVE -  
DETECTS LIGHT**




**OPTIC NERVE -  
CN II VISION**

**CN III - OCULOMOTOR - parasympathetics  
from Ciliary Ganglion in Short Ciliary nerves**



**CONTRACTOR PUPIL-  
CIRCULAR SMOOTH MUSCLE;  
PARASYMPATHETICS - CN III**



# PUPILLARY LIGHT REFLEX

**CN II - OPTIC NERVE - DETECTS LIGHT**

**CN III - OCULOMOTOR - parasympathetics from Ciliary Ganglion in Short Ciliary nerves**

**CN III - OCULOMOTOR**


**CN III - OCULOMOTOR**

**CN II - OPTIC NERVE**

**CONSENSUAL REFLEX – PUPIL CONSTRICTS IN OPPOSITE EYE**

**DIRECT REFLEX – PUPIL CONSTRICTS IN SAME EYE**

**SHINE LIGHT**



## 2. CORNEAL REFLEX - V TO VII

AFFERENT ARM OF REFLEX

**SENSORY  
STIMULUS**

**TOUCH  
CORNEA**

**TRIGEMINAL -  
V1 - LONG  
CILIARY NERVES  
TO CORNEA**



EFFERENT ARM OF REFLEX

**MOTOR  
RESPONSE**

**CLOSE  
EYELID**


**FACIAL -  
VII - MOTOR TO  
ORBICULARIS  
OCULI (SVE)**




# CORNEAL REFLEX - V to VII



**V - TOUCH CORNEA**



**VII - CLOSE EYELID**



**ORBICULARIS OCULI M.**

**LONG CILIARY NERVES (V1) - SOMATIC SENSORY TO CORNEA**

**- Palpebral part - Close eyelids  
- Orbital part - Buries eyelids, Ex. sandstorm  
BRANCHIOMOTOR - VII**

**SHORT CILIARY NERVES (III),  
CILIARY GANGLION  
PARASYMPATHETIC**

### 3. GAG REFLEX - IX to X \* \*

AFFERENT ARM OF REFLEX

**SENSORY  
STIMULUS**

**TOUCH  
ORO-  
PHARYNX**

EFFERENT ARM OF REFLEX

**MOTOR  
RESPONSE**

**PATIENT GAGS -  
CONTRACT  
PHARYNGEAL  
MUSCLES**



# CRANIAL NERVES LECTURE

## GAG REFLEX



### IX - SENSORY INNERVATION TO OROPHARYNX

X - INNERVATES ALL MUSCLES OF PHARYNX (except Stylopharyngeus)

All Pharynx is Visceral Sensory In 3 Cranial Nerves

SOMATIC SENSORY

VISCERAL SENSORY




IX AND X - LEAVE MEDULLA, EXIT BY JUGULAR FORAMEN - CAN DIAGNOSE DAMAGE IN BRAINSTEM BY TESTING REFLEXES

# 4. STRETCH REFLEX OF MUSCLES OF MASTICATION



- **JAW JERK REFLEX** - sensory and motor in Trigeminal V3

## STRETCH REFLEX




## STRETCH REFLEX IN MUSCLES OF MASTICATION

### TAP DOWN ON CHIN



MEDIAL PTERYGOID

**STRETCH MUSCLES THAT CLOSE MOUTH (ELEVATE MANDIBLE) TEMPORALIS**



MASSETER

**TEST FUNCTION OF TRIGEMINAL NERVE (V3)**

# SUMMARY OF SPINAL, CRANIAL NERVE REFLEXES

## SPINAL REFLEXES AND DISORDERS

REFLEX	STIMULUS/SENSE ORGAN(S) EXCITED	NORMAL RESPONSE	UPPER MOTOR NEURON DISORDERS
Stretch (Myotatic, Deep Tendon) Reflex – Compensatory maintain position (ex. riding on moving bus)	Rapid Stretch of muscle (test: tap on muscle tendon) Excites Muscle Spindle Primary (Ia) and Secondary (II) sensory neurons (NOT Golgi Tendon Organ)	Stretched muscle contracts rapidly (monosynaptic connection); also Excite synergist and Inhibit antagonist Note: Gamma motor neurons can enhance stretch reflexes, tell patient to relax before test	<u>Hyperreflexia</u> - (increase) - characteristic of Upper Motor Neuron lesions (ex. spinal cord injury, damage Corticospinal tract); note: <u>Clonus</u> = hyperreflexia with repetitive or sustained contractions to single stimulus
Autogenic Inhibition - Limits Muscle Tension	Large force on tendon excites Golgi Tendon Organ Ib (test: pull on muscle when resisted)	Muscle tension decreases; Also inhibit synergist muscles; excite antagonist muscles	<u>Clasped Knife Reflex</u> - occurs in Upper Motor Neuron lesions - forceful stretch of muscle is first resisted then collapses
Flexor Reflex - Protective avoidance reflex	Sharp, painful stimulus, as in stepping on nail; Excites - Cutaneous and pain receptors (test: stroke foot with pointed object)	Limb is rapidly withdrawn from stimulus; protective reflex; also inhibit extensors of same limb and excite extensors of opposite limb (Crossed Extensor Reflex)	<u>Babinski sign</u> -toes extend (dorsiflex) to cutaneous stimulus of sole of foot (normally plantar flex); characteristic of Upper Motor Neuron lesion

## REFLEXES OF CRANIAL NERVES

REFLEX	STIMULUS	SENSORY	RESPONSE	CLINICAL
Pupillary Light Reflex (II to III)	Test: Shine light in eye	Light detected by Optic Nerve	Excite Constrictor of pupil of eye (III Short Ciliary nerves (Ciliary Ganglion, parasympathetic)	Extensively used to check CN II; Absence of Pupillary Light Reflex can indicate catastrophe (brain herniation)
Corneal Reflex (V to VII)	Touch cornea of eye with cotton	Touch detected by Long Ciliary nerves (V1), Somatic sensory	Close eye (VII to Orbicularis Oculi muscle) Branchiomotor	Absence of Corneal Reflex; Test for damage to V1 sensory, VII motor
Gag Reflex (IX to X)	Test: Touch posterior tongue, oropharynx;	Excites Visceral Sensory endings in Glossopharyngeal N. (IX)	Excite muscles of pharynx, palate; Vagus N. (X), Branchiomotor	Other symptoms of Vagus damage (X); Patient Say's Ahh: soft palate not elevated on ipsilateral side (paralyze Levator Palati); uvula deviated away from side of lesion
Jaw Jerk Reflex Stretch (Deep Tendon) Reflex (V to V)	Test: tap down on mandible; Stretch muscles of mastication (ex. Masseter)	Excites Muscle Spindle sensory neurons in Trigeminal nerve (V)	Contract muscles that elevate mandible Motor - V3	<u>Hyporeflexia</u> - indicates Trigeminal nerve damage